



Decoding primordial physics through gravitational waves

L. Sriramkumar

Centre for Strings, Gravitation and Cosmology, Department of Physics,
Indian Institute of Technology Madras, Chennai

J. V. Narlikar Memorial Conference on Cosmology and Astrophysics
Department of Physics, Visva-Bharati, Santiniketan
November 3–4, 2025



Books that inspired me as an undergraduate and master's student



Books by Prof. Jayant Narlikar that drew me to astronomy and astrophysics as an undergraduate and master's student.



Books that I studied as a master's and doctoral student



Books by Prof. Jayant Narlikar that I studied as a master's and doctoral student.



Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 GWs provide a new window to the universe
- 4 Generation of GWs in the early universe
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook




Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 GWs provide a new window to the universe
- 4 Generation of GWs in the early universe
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook



Timeline of the universe




► Observations of GWs

A pictorial timeline of the universe, according to the standard model of cosmology¹.

¹See http://wmap.gsfc.nasa.gov/media/060915/060915_CMB_Timeline150.jpg.





Horizon problem



The radiation from the CMB arriving at us from regions separated by more than the Hubble radius at the surface of last scattering, which subtends an angle of about 1° today, could not have interacted before decoupling.




Resolution of the horizon problem in the inflationary scenario



Another illustration of the horizon problem (on the left), and an illustration of its resolution (on the right) through an early and sufficiently long epoch of inflation².

²Images from W. Kinney, [astro-ph/0301448](https://arxiv.org/abs/astro-ph/0301448).

Time and duration of inflation





Inflation—a brief period of accelerated expansion—is expected to have taken place during the very early stages of the universe³.

³Image from P. J. Steinhardt, *Sci. Am.* **304**, 18 (2011).



Inflationary attractor



► Back to USR inflation

Evolution of the scalar field in the popular Starobinsky model, which leads to slow roll inflation, is indicated (as circles, in blue and red) at regular intervals of time (on the left). Illustration of the behavior of the scalar field in phase space (on the right)⁴.

⁴Figure from H. V. Ragavendra, *Observational imprints of non-trivial inflationary dynamics over large and small scales*, Ph.D. Thesis, Indian Institute of Technology Madras, Chennai, India (2022).



Origin of the primordial perturbations

Scalar perturbations:

- ◆ The quantum fluctuations associated with the scalar fields that drive inflation are responsible for the primordial perturbations.
- ◆ The scalar perturbations leave the largest imprints on the CMB, and are primarily responsible for the inhomogeneities in the distribution of matter in the universe.


Tensor perturbations:

- ◆ The tensor perturbations, i.e. gravitational waves (GWs), can be generated even in the absence of sources.
- ◆ GWs are small disturbances in a given spacetime (much like ripples in water), which travel at the speed of light. They satisfy the wave equation in the given background spacetime.
- ◆ GWs are transverse in nature and are characterized by two degrees of polarization⁵.

⁵See, for example, J. B. Hartle, *Gravity: An Introduction to Einstein's General Relativity* (Pearson Education, Delhi, 2003).



Behavior of the comoving wave number and Hubble radius



► Back to reheating

► Back to formation of PBHs

► Back to SGWs and PTA data

Behavior of the comoving wave number k (horizontal lines in different colors) and the comoving Hubble radius $d_H/a = (aH)^{-1}$ (in green) across different epochs⁶.

⁶Md. R. Haque, D. Maity, T. Paul and L. Sriramkumar, Phys. Rev. D **104**, 063513 (2021).




Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 GWs provide a new window to the universe
- 4 Generation of GWs in the early universe
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook



CMB angular power spectrum from Planck




The CMB TT angular power spectrum from the Planck 2018 data (red dots with error bars) and the best fit Λ CDM model with a power law primordial spectrum (solid blue curve)⁷.

⁷Planck Collaboration (N. Aghanim *et al.*), *Astron. Astrophys.* **641**, A6 (2020).



Performance of inflationary models in the n_s - r plane




Latest constraints on n_s and r from ACT, in combination with other data sets, compared to the theoretical predictions of some of the popular inflationary models⁸.

⁸ACT Collaboration (E. Calabrese *et al.*), arXiv:2503.14454 [astro-ph.CO].



Prospects of observing the imprints of primary GWs in the CMB



The B-mode angular power spectra of the CMB resulting from the primary tensor perturbations for models with $r_{0.05} = 5 \times 10^{-2}$ and $r_{0.05} = 4 \times 10^{-3}$ have been plotted, along with the CMB lensing signal and the instrumental noise of a LiteBIRD-like configuration⁹.

⁹D. Paoletti, F. Finelli, J. Valiviita and M. Hazumi, Phys. Rev. D **106**, 083528 (2022).




Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 **GWs provide a new window to the universe**
- 4 Generation of GWs in the early universe
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook




Sources and spectral range of GWs



Different sources of GWs and the corresponding detectors¹⁰.

¹⁰ J. B. Hartle, *Gravity: An Introduction to Einstein's General Relativity* (Pearson Education, Delhi, 2003).

Probing the primordial universe through GWs



GWs provide a unique window to probe the primordial universe¹¹.

¹¹ Image from <https://gwpo.nao.ac.jp/en/gallery/000061.html>.




Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 GWs provide a new window to the universe
- 4 Generation of GWs in the early universe**
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook




Evolution of the scalar field in an inflationary potential



The evolution of the scalar field in the so-called Starobinsky model has been indicated (as circles, in blue and red) at regular intervals of time. Inflation is terminated as the field approaches the bottom of the potential (near the light blue dot). Thereafter, the field oscillates at the bottom of the potential (indicated by the red dots).



Effects on primary GWs due to late time entropy production




► Evolution of comoving lengths

The dimensionless spectral energy density of primary GWs observed today $\Omega_{\text{GW}}(f)$ has been plotted in a scenario involving late time production of entropy¹².

¹²Md. R. Haque, D. Maity, T. Paul and L. Sriramkumar, Phys. Rev. D **104**, 063513 (2021).



Coalescence of compact binaries observed by LIGO




The third GW Transient Catalog of mergers involving BHs and neutron stars observed by the LIGO-Virgo-KAGRA collaboration¹³.

¹³Image from <https://www.ligo.caltech.edu/LA/image/ligo20211107a>.




Formation of BHs in the early universe



BHs can form when perturbations with significant amplitudes reenter the Hubble radius during the radiation dominated epoch¹⁴.

¹⁴Figure from G. Franciolini, [arXiv:2110.06815 \[astro-ph.CO\]](https://arxiv.org/abs/2110.06815).

Amplitude for producing significant number of primordial BHs (PBHs)





In order to form significant number of PBHs, the amplitude of the perturbations on small scales has to be large enough such that the dimensionless amplitude of the scalar perturbation is close to unity¹⁵.

¹⁵Figure credit G. Franciolini.



Single-field models admitting ultra slow roll (USR) inflation




► Inflationary attractor

Potentials which contain a point of inflection generically admit a period of USR inflation¹⁶.

¹⁶See, for example, C. Germani and T. Prokopec, *Phys. Dark Univ.* **18**, 6 (2017);
I. Dalianis, A. Kehagias and G. Tringas, *JCAP* **01**, 037 (2019).

Figures credits, H. V. Ragavendra and S. Maity.

Power spectra in models permitting USR inflation




[Back to analytical form of the power spectrum](#)

Scalar (in red) and the tensor (in blue) power spectra arising in different single-field models that permit a period of USR inflation¹⁷.

¹⁷ H. V. Ragavendra, P. Saha, L. Sriramkumar and J. Silk, Phys. Rev. D **103**, 083510 (2021);

Also see H. V. Ragavendra and L. Sriramkumar, Galaxies **11**, 34 (2023).

Formation of PBHs in single-field and two-field models of inflation




The fraction of PBHs contributing to the dark matter density today $f_{\text{PBH}}(M)$ arising in different single-field¹⁸ and two-field¹⁹ inflationary models.

¹⁸ H. V. Ragavendra, P. Saha, L. Sriramkumar and J. Silk, Phys. Rev. D **103**, 083510 (2021).

¹⁹ M. Braglia, D. K. Hazra, F. Finelli, G. F. Smoot, L. Sriramkumar and A. A. Starobinsky, JCAP **08**, 001 (2020).



Secondary GWs in the two-field model



The dimensionless density parameter $\Omega_{GW}(f)$ arising in the two-field model has been plotted for a set of initial conditions for the background fields as well as a range of values of a parameter describing the model²⁰.


²⁰M. Braglia, D. K. Hazra, F. Finelli, G. F. Smoot, L. Sriramkumar and A. A. Starobinsky, JCAP **08**, 001 (2020).

Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 GWs provide a new window to the universe
- 4 Generation of GWs in the early universe
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook



Pulsar timing arrays (PTAs)




The PTAs monitor an array of millisecond pulsars²¹.

²¹See https://ipta.github.io/mock_data_challenge/.



Stochastic GW background observed by pulsar timing arrays (PTAs)



The Bayesian evidence for a variety of astrophysical and cosmological sources for the stochastic GW background suggested by the observations of the PTAs²².

²² NANOGrav Collaboration (G. Agazie *et al.*), *Astrophys. J. Lett.* **951**, 1 (2023).



Shape of the inflationary scalar power spectrum

We assume that the inflationary scalar power spectrum is given by²³

$$\mathcal{P}_s(k) = A_s \left(\frac{k}{k_*} \right)^{n_s-1} + A_0 \begin{cases} \left(\frac{k}{k_{\text{peak}}} \right)^4 & k \leq k_{\text{peak}}, \\ \left(\frac{k}{k_{\text{peak}}} \right)^{n_0} & k \geq k_{\text{peak}}, \end{cases}$$

where A_s and n_s are the amplitude and spectral index of the power spectrum at the CMB pivot scale of $k_* = 0.05 \text{ Mpc}^{-1}$.

► Power spectra in USR inflation

We set the reheating temperature to the rather low value of $T_{\text{re}} = 50 \text{ MeV}$.

We shall assume that the threshold value of the density contrast for the formation of PBHs is given by²⁴:


$$\delta_c^{\text{an}} = \frac{3(1+w_{\text{re}})}{5+3w_{\text{re}}} \sin^2 \left(\frac{\pi \sqrt{w_{\text{re}}}}{1+3w_{\text{re}}} \right).$$

²³For other forms of spectra, see G. Domènech, S. Pi, A. Wang and J. Wang, arXiv:2402.18965 [astro-ph.CO].

²⁴In this context, see T. Harada, C.-M. Yoo, and K. Kohri, Phys. Rev. D **88**, 084051 (2013).



Generation of secondary GWs during the epoch of reheating





The dimensionless spectral energy density of the secondary GWs today $\Omega_{\text{GW}}(f)$ is plotted for a given reheating temperature and the best-fit values of the parameters in the different models²⁵.

²⁵S. Maity, N. Bhaumik, Md. R. Haque, D. Maity and L. Sriramkumar, JCAP **01**, 118 (2025).




Power spectra and the extent of PBHs formed



Scalar power spectra (on the left) and the extent of PBHs formed (on the right). We have assumed a specific reheating temperature and have plotted the fraction of PBHs that constitute the dark matter density today, viz. $f_{\text{PBH}}(M)$, for the best-fit values of the parameters in the different models²⁶.

²⁶S. Maity, N. Bhaumik, Md. R. Haque, D. Maity and L. Sriramkumar, JCAP **01**, 118 (2025).

Constraints on the epoch of reheating



Constraints on the parameters of the two models R3pB (on the left) and R2pB (on the right), arrived at upon comparison with the NANOGrav 15-year data²⁷.

²⁷ S. Maity, N. Bhaumik, Md. R. Haque, D. Maity and L. Sriramkumar, JCAP **01**, 118 (2025).



Plan of the talk

- 1 Inflationary scenario
- 2 Constraints on inflation from the CMB data
- 3 GWs provide a new window to the universe
- 4 Generation of GWs in the early universe
- 5 Observations by the PTAs and the stochastic GW background
- 6 Outlook



Outlook

- ◆ The increasingly precise observations of the CMB by future missions such as Lite-BIRD (Light satellite for the studies of B-mode polarization and Inflation from cosmic background Radiation Detection), Primordial Inflation Explorer (PIXIE) and Exploring Cosmic History and Origin (ECHO, a proposed Indian effort) can be expected to help us improve the current constraints on the primordial correlations.
- ◆ The observations by LIGO are a culmination of almost fifty years of effort to detect GWs. They have opened up a new window to observe the universe.
- ◆ The observations by the PTAs and their possible implications for the stochastic GW background offer a wonderful opportunity to understand the physics operating in the early universe.
- ◆ Over the coming decades, GW observatories such as the Laser Interferometer Space Antenna, Einstein Telescope and Cosmic Explorer, can be expected to provide us with an unhindered view of the primordial universe.



This talk was based on...

- ◆ M. Braglia, D. K. Hazra, F. Finelli, G. F. Smoot, L. Sriramkumar and A. A. Starobinsky, *Generating PBHs and small-scale GWs in two-field models of inflation*, JCAP **08**, 001 (2020) [arXiv:2005.02895 [astro-ph.CO]].
- ◆ H. V. Ragavendra, P. Saha, L. Sriramkumar and J. Silk, *Primordial black holes and secondary gravitational waves from ultra slow roll and punctuated inflation*, Phys. Rev. D **103**, 083510 (2021) [arXiv:2008.12202 [astro-ph.CO]].
- ◆ Md. R. Haque, D. Maity, T. Paul and L. Sriramkumar, *Decoding the phases of early and late time reheating through imprints on primordial gravitational waves*, Phys. Rev. D **104**, 063513 (2021) [arXiv:2105.09242 [astro-ph.CO]].
- ◆ H. V. Ragavendra and L. Sriramkumar, *Observational imprints of enhanced scalar power on small scales in ultra slow roll inflation and associated non-Gaussianities*, Galaxies **11**, 34 (2023) [arXiv:2301.08887 [astro-ph.CO]].
- ◆ S. Maity, N. Bhaumik, Md. R. Haque, D. Maity and L. Sriramkumar, *Constraining the history of reheating with the NANOGrav 15-year data*, JCAP **01**, 118 (2025) [arXiv:2403.16963 [astro-ph.CO]].



Collaborators I



Matteo Braglia



Dhiraj Hazra



Fabio Finelli



George Smoot



Alexei Starobinsky



H. V. Ragavendra



Pankaj Saha



Joseph Silk

Collaborators II



Md. Riajul Haque



Debaprasad Maity



Tanmoy Paul



Suvashis Maity



Nilanjandev Bhaumik



Thank you for your attention