

Introduction to cosmology

L. Sriramkumar

Centre for Strings, Gravitation and Cosmology, Department of Physics,
Indian Institute of Technology Madras, Chennai

Summer Training Programme in Physics
Department of Materials Science, University of Madras, Chennai
Science City, Department of Higher Education, Government of Tamil Nadu
June 30, 2023

Plan of the talk

1 A survey of the universe



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology
- 6 Useful references




Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology
- 6 Useful references



An infrared image of our galaxy




Our galaxy – the Milky Way – as observed by the COsmic Background Explorer (COBE) satellite at the infrared wavelengths¹. The diameter of the disc of our galaxy is, approximately, 45×10^3 ly or 15 kpc (*i.e.* a kilo parsec, with $1 \text{ pc} \simeq 3.26 \text{ ly}$). It contains about 10^{11} stars such as the Sun, and its mass is about $2 \times 10^{12} M_{\odot}$.

¹Image from http://aether.lbl.gov/www/projects/cobe/cobe_pics.html.



Stellar parallax²



The baseline of the earth's orbit of **2 AU** can be used to determine the distances of nearby stars through trigonometric parallax.

²Image from <http://find.uchicago.edu/~pryke/compton/slides2/mgp00007.html>.



The parsec

- ◆ **Parsec (pc)**: The distance to an object whose parallax is $1''$ due to the baseline of the earth's orbit of 2 AU .

From the figure in the previous slide, it is clear that

$$d = (1 \text{ AU} / \tan p) \simeq (1/p) \text{ AU},$$

where we have assumed that the angle p is small. If p is expressed in units of *arcseconds*, we find that

$$d = \left(\frac{2.063 \times 10^5}{p''} \right) \text{ AU}.$$

Note that for $p = 1''$, $d = 2.063 \times 10^5 \text{ AU} = 1 \text{ pc} = 3.26 \text{ ly} = 3.0857 \times 10^{16} \text{ m}$.



Our galactic neighbors and the local group³






Left: The Andromeda galaxy and its two companion galaxies. The Andromeda galaxy is very similar to our galaxy and is located at a distance of about **700** kpc.

Right: The Triangulum galaxy M33. These galaxies, along with our galaxy, are major members of a local group of about **30** galaxies that are bound gravitationally. The size of the local group is estimated to be about **1.3** Mpc.

³Images from <http://www.seds.org/messier/m/m031.html> and <http://www.seds.org/messier/m/m033.html>.



Varieties of galaxies⁴



Left: The disk galaxy NGC 4565 seen edge on in this image from the Sloan Digital Sky Survey (SDSS). The galaxy has a clear bulge, but little light can be seen from its halo.



Center: An image of the spiral galaxy NGC 3187 from SDSS.

Right: CGCG 180-023 is a superb example of a ring galaxy. Ring galaxies are believed to form when a compact smaller galaxy plunges through the center of a larger more diffuse rotating disk galaxy.

⁴Images from <http://www.sdss.org/iotw/archive.html> and <http://cosmo.nyu.edu/hogg/rc3>.



The Virgo, the Coma and the Hercules cluster of galaxies⁵



Left: The Virgo cluster, whose center is considered to be located at a distance of about 20 Mpc. Consisting of over 100 galaxies, it strongly influences the nearby galaxies and galaxy groups gravitationally due to its enormous mass.


Center: The Coma cluster of galaxies, which contains more than 1000 bright galaxies. It is about 20 Mpc across, and is located at a distance of about 100 Mpc.

Right: An SDSS image of the Hercules galaxy cluster that is located at a distance of about 100 Mpc from us.

⁵ Images from <http://apod.nasa.gov/apod/ap000220.html>, <http://www.astr.ua.edu/gifimages/coma.html> and <http://www.sdss.org/iotw/archive.html>.



Deepest views in space




An ultra deep field image from the Hubble Space Telescope (HST). The image contains a bewildering variety of galaxy shapes and colors⁶.

⁶Image from <http://hubblesite.org/newscenter/archive/releases/2014/27>.



Surveying the universe




A schematic drawing showing the directions of the regions observed by the 2 degree field (2dF) redshift survey with respect to our galaxy⁷. The survey regions actually extend more than 10^5 times further than shown here.

⁷Image from <http://magnum.anu.edu.au/~TDFgg/Public/Pics/2dF3D.jpg>.



Distribution of galaxies in the universe



The distribution of more than two million galaxies as observed by the 2dF redshift survey⁸. (Note that each dot in the picture represents a galaxy.) The density and the 'radius' of the universe are estimated to be about 10^{-28} kg/m³ and 3000 Mpc, respectively.

⁸Image from http://magnum.anu.edu.au/~TDFgg/Public/Pics/2dFGRS_top_view.gif.



The Sloan digital sky survey

- ◆ The Sloan Digital Sky Survey (SDSS) is one of the most ambitious and influential surveys in the history of astronomy⁹.
- ◆ Over eight years of operations, it has obtained deep, multi-color images covering more than a quarter of the sky and created three-dimensional maps containing more than 930,000 galaxies and more than 120,000 quasars.

► Play SDSS movie

⁹See, <http://www.sdss.org/>.



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology
- 6 Useful references




Continuous, emission and absorption spectra¹⁰

A typical continuous spectrum from an opaque hot body:



Emission spectrum, as from a given element:




Absorption spectrum, as due to an intervening cool gas:



¹⁰Images from <http://hea-www.harvard.edu/~efortin/thesis/html/Spectroscopy.shtml>.

Typical spectra of galaxies¹¹



Spectra of some spiral galaxies. The spectra usually contain characteristic emission and absorption lines.

¹¹ Image from <http://astronomy.nmsu.edu/nicole/teaching/ASTR505/lectures/lecture26/slide01.html>.

The 'Doppler effect' and redshift¹²

If the source is receding, the spectrum will be red-shifted



when compared to the spectrum in the source's frame



The redshift z of the receding source is defined as:


$$1 + z = \frac{\lambda_O}{\lambda_E} = \frac{\omega_E}{\omega_O},$$

where λ_O and ω_O denote the observed wavelength and frequency of the source, while λ_E and ω_E denote its emitted wavelength and frequency, respectively.

¹²Images from <http://www.astronomynotes.com/light/s10.htm>.



Runaway galaxies¹³





Spectra of four different galaxies from the **2dF redshift survey**. On top left is the spectrum of a star from our galaxy, while on the bottom right we have the spectrum of a galaxy that has a redshift of $z = 0.246$. The other two galaxies show prominent H α emission lines, which have been redshifted from the rest frame value of **6563 Å**.

¹³Image from http://outreach.atnf.csiro.au/education/senior/astrophysics/spectra_astro_types.html.



Relation between the velocity and the distance of galaxies¹⁴



Left: The original Hubble data. The slope of the two fitted lines are about 500 km/sec/Mpc and 530 km/sec/Mpc.

Right: A more recent Hubble diagram. The slope of the straight line is about 72 km/sec/Mpc. The small red region in the lower left marks the span of Hubble's original diagram.

¹⁴R. Kirshner, Proc. Natl. Acad. Sci. USA **101**, 8 (2004).




The Friedmann-Lemaître-Robertson-Walker metric

The homogeneous, isotropic and expanding universe can be described by the following Friedmann-Lemaître-Robertson-Walker (FLRW) line element:


$$ds^2 = dt^2 - a^2(t) \left[\frac{dr^2}{(1 - \kappa r^2)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right],$$

where t is the cosmic time and $a(t)$ denotes the scale factor, while $\kappa = 0, \pm 1$.


The quantity κ denotes the spatial geometry of the universe. It can be flat ($\kappa = 0$), closed ($\kappa = 1$) or open ($\kappa = -1$) depending on the total energy density of matter present in the universe¹⁵.



Positive Curvature



Negative Curvature



Flat Curvature

¹⁵Image from http://abyss.uoregon.edu/~js/lectures/cosmo_101.html.



The Friedmann equations

If ρ and p denote the energy density and pressure of the smooth component of the matter field that is driving the expansion, then the Einstein's equations for the FLRW metric lead to the following equations for the scale factor $a(t)$:

$$H^2 + \frac{\kappa}{a^2} = \frac{8\pi G}{3} \rho$$


and

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} (\rho + 3p),$$

where $H = \dot{a}/a$ is the Hubble parameter.



Visualizing the expanding universe



A two-dimensional analogy for the expanding universe¹⁶. The yellow blobs on the expanding balloon denote the galaxies. Note that the galaxies themselves do not grow, but the distance between the galaxies grows and the wavelengths of the photons shift from blue to red as the universe expands.

¹⁶Image from <http://www.astro.ucla.edu/~wright/balloon0.html>.



The cosmological redshift

Recall that, we had defined the redshift z of a receding source as follows:

$$1 + z = \frac{\omega_E}{\omega_O},$$

where ω_O and ω_E denote the observed and emitted frequencies, respectively.

In an expanding universe, it can be shown that the frequency of electromagnetic radiation decreases with the expansion as follows:

$$\omega(t) \propto \frac{1}{a(t)},$$

where $a(t)$ is the scale factor that characterizes the expansion.

Therefore, in terms of the scale factor, the cosmological redshift z is given by

$$\frac{a_0}{a(t)} = 1 + z,$$

where a_0 denotes the value of the scale factor *today* (i.e. at $t = t_0$).



The cosmological parameters

In terms of the redshift z , the first of the Friedmann equations can be written as

$$\left[\frac{H(z)}{H_0} \right]^2 = \Omega_{\text{NR}} (1+z)^3 + \Omega_{\text{R}} (1+z)^4 + \Omega_{\Lambda} - (\Omega - 1) (1+z)^2,$$

where $H_0 \equiv (\dot{a}/a)_{t=t_0}$ is the Hubble constant, $\Omega_i = \rho_i/\rho_{\text{C}}$ with ρ_{C} being the critical density given by



$$\rho_{\text{C}} = \frac{3 H_0^2}{8 \pi G}$$

and $\Omega = \Omega_{\text{NR}} + \Omega_{\text{R}} + \Omega_{\Lambda}$.

The quantities H_0 , Ω_{NR} , Ω_{R} and Ω_{Λ} are four of the cosmological parameters that are to be determined by observations.



Discovery of the cosmic microwave background (CMB)




The horn antenna used by Penzias and Wilson (on the left) and the CMB as observed by them (on the right)¹⁷.

► Anisotropies in the CMB

¹⁷In this context, see, for instance, S. G. Brush, *Sci. Am.* **267**, 62 (1992).



Complete spectrum of the cosmological background radiation




The energy density spectrum of cosmological background radiation has been plotted as a function of wavelength¹⁸. Note that the CMB contributes the most to the overall background radiation.

¹⁸Figure from, D. Scott, [arXiv:astro-ph/9912038](https://arxiv.org/abs/astro-ph/9912038).



Spectrum of the CMB



The spectrum of the CMB as measured by the COBE satellite¹⁹. It is a perfect Planck spectrum (corresponding to a temperature of 2.725° K) which is unlikely to be bettered in the laboratory. The error bars have been amplified 400 times so that they are visible!

¹⁹Image from http://www.astro.ucla.edu/~wright/cosmo_01.htm.



The radiation and the matter dominated epochs

In an evolving universe, the temperature of the CMB goes as

$$T \propto \frac{1}{a(t)},$$

so that the energy density of radiation behaves as

$$\rho_{\text{R}} \propto \frac{1}{a^4(t)}.$$

In contrast, the energy density of non-relativistic (*i.e.* pressureless) matter goes as

$$\rho_{\text{NR}} \propto \frac{1}{a^3(t)}.$$


Observations indicate that, today,

$$\rho_{\text{R}} \simeq \frac{\rho_{\text{NR}}}{10^4}.$$

This points to the fact that matter and radiation would have interacted strongly and, hence, would have been in thermal equilibrium, when the universe was about 10^4 times smaller.



The big bang model seems popular!




The current view of the universe, encapsulated in the hot big bang model, seems popular. The above image is a screen grab from the theme song of the recent American sitcom 'The Big Bang Theory'²⁰!

²⁰See http://www.cbs.com/shows/big_bang_theory/.



Decoupling of matter and radiation





The CMB arises because matter and radiation cease to interact at an early time²¹.

²¹Image from W. H. Kinney, [arXiv:astro-ph/0301448v2](https://arxiv.org/abs/astro-ph/0301448v2).



The last scattering surface and the freestreaming CMB photons




The CMB photons streams to us freely from the last scattering surface when radiation decoupled from matter²².

²²Image from <http://planck.caltech.edu/epo/epo-cmbDiscovery4.html>.



Projecting the last scattering surface




The temperature of the CMB on the last scattering surface can be projected on to a plane as the surface of the Earth is often projected²³.

²³Image from <http://hyperphysics.phy-astr.gsu.edu/hbase/Astro/planckcmb.html>.



The extent of isotropy of the CMB




The fluctuations in the temperature of the CMB as seen by COBE²⁴. The CMB turns out to be isotropic to one part in 10^5 .

► The smooth CMB

²⁴Image from http://aether.lbl.gov/www/projects/cobe/COBE_Home/DMR_Images.html.



The abundance of light elements – Theory




The relative abundances of the light elements in the early radiation dominated epoch have been plotted as a function of temperature²⁵.

²⁵Image from <http://www.astro.ucla.edu/~wright/BBNS.html>.



Abundance of light elements – Observations²⁶




The graph to the left contains the theoretically predicted abundance versus the density for the light elements as curves, the observed abundances as horizontal stripes and the derived baryon density as the vertical stripe. Note that a single value of the baryon density fits all the four abundances, and it is found that $\Omega_B h^2 \simeq 0.022$, where $H_0 = 100 h$ km/sec/Mpc.

²⁶Image from <http://www.astro.ucla.edu/~wright/BBNS.html>.



Rotation curves of spiral galaxies




The observed rotation curve of the spiral galaxy M33 (yellow and blue points with error bars), and the predicted curve from the distribution of visible matter (gray line). The observed curve can be accounted for by embedding the galaxy in a dark matter halo²⁷.

²⁷Image from https://en.wikipedia.org/wiki/Galaxy_rotation_curve.



Gravitational lensing reveals the distribution of matter



A near perfect Einstein ring! The ring is formed due to the gravitational field of the foreground luminous red galaxy which distorts the light from a more distant blue galaxy²⁸.

²⁸Image from <https://apod.nasa.gov/apod/ap111221.html>.



Supernovae can be as bright as the host galaxy²⁹



Supernova 1994D, visible as the bright spot on the lower left, occurred in the outskirts of disk galaxy NGC 4526.

²⁹Image from <http://apod.nasa.gov/apod/ap981230.html>.



A supernova explosion in a distant galaxy³⁰




Left: A supernova at the redshift of 0.28 caught at maximum light by the Supernova Legacy Survey (SNLS).

Right: The supernova after it has faded.

³⁰Images from C. J. Pritchett *et. al.*, [arXiv:astro-ph/0406242v1](https://arxiv.org/abs/astro-ph/0406242v1).



Supernovae data and the need for a cosmological constant




The luminosity distance $H_0 d_L$ plotted as a function of the redshift z for spatially flat cosmological models³¹. The black points are from the ‘Gold’ data sets and the red points are the data from the Hubble Space Telescope³².

³¹Figure from T. R. Choudhury and T. Padmanabhan, *Astron. Astrophys.* **429**, 807 (2005).

³²R. A. Knop *et. al.*, *Astrophys. J.* **598**, 102 (2003); A. G. Riess *et. al.*, *Astrophys. J.* **607**, 665 (2004).



Joint constraints on Ω_{NR} and Ω_{Λ} ³³




Joint constraints on Ω_{NR} and Ω_{Λ} from the observations of supernovae, CMB and galaxy clustering. Note that a cosmology with $\Omega_{NR} = 1$ and $\Omega_{\Lambda} = 0$ is ruled out to 99% confidence level, while a universe with $\Omega_{NR} \simeq 0.3$ and $\Omega_{\Lambda} \simeq 0.7$ proves to be a good fit to the data. The figure also contains the constraints that can be expected from the planned Supernova/Acceleration Probe (SNAP).

³³Figure from G. Aldering *et. al.*, arXiv:astro-ph/0209550v1.



Matter content of the universe



A pie chart of the matter content of the universe today³⁴.

³⁴Image from http://map.gsfc.nasa.gov/media/060916/060916_UniversePie300.jpg.




Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology
- 6 Useful references




The horizon problem



The radiation from the CMB arriving at us from regions separated by more than the Hubble radius at the last scattering surface, which subtends an angle of about 1° today, could not have interacted before decoupling.



Inflation resolves the horizon problem




An early and sufficiently long epoch of inflation resolves the horizon problem³⁵.

³⁵Image from W. H. Kinney, [arXiv:astro-ph/0301448v2](https://arxiv.org/abs/astro-ph/0301448v2).



Bringing the modes inside the Hubble radius




Behavior of the physical wavelength $\lambda_P \propto a$ (in blue) and the Hubble radius $d_H = H^{-1}$ (in red) in the inflationary scenario³⁶. Note that the scale factor is expressed in terms of e-folds N as $a(N) \propto e^N$.

³⁶See, for example, E. W. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley Publishing Company, New York, 1990), Fig. 8.4.



The time and duration of inflation



Inflation – a brief period of accelerated expansion – is expected to have taken place during the very stages of the universe³⁷.

³⁷Image from P. J. Steinhardt, *Sci. Am.* **304**, 18 (2011).



Necessary conditions for inflation³⁸

If we require that $\lambda_P < d_H$ at a sufficiently early time, then we need to have an epoch wherein λ_P decreases faster than the Hubble scale *as we go back in time*, i.e. a regime during which

$$-\frac{d}{dt} \left(\frac{\lambda_P}{d_H} \right) < 0 \quad \rightarrow \quad \ddot{a} > 0.$$

From the Friedmann equations, we have

$$\left(\frac{\ddot{a}}{a} \right) = - \left(\frac{4\pi G}{3} \right) (\rho + 3p).$$

Evidently, $\ddot{a} > 0$ when

$$(\rho + 3p) < 0.$$

³⁸See, for example, L. Sriramkumar, Curr. Sci. **97**, 868 (2009).



Scalar fields can drive inflation³⁹

In a smooth Friedmann universe, the energy density ρ and pressure p corresponding to a homogeneous scalar field, say, ϕ are given by

$$\rho = \left(\dot{\phi}^2/2 \right) + V(\phi) \quad \text{and} \quad p = \left(\dot{\phi}^2/2 \right) - V(\phi),$$

and the scalar field satisfies the following equation of motion:

$$\ddot{\phi} + 3H\dot{\phi} + (dV/d\phi) = 0.$$

In such a case, the condition $(\rho + 3p) < 0$ simplifies to


$$\dot{\phi}^2 < V(\phi).$$

This condition can be achieved if the scalar field ϕ is initially displaced from a minima of the potential, and inflation will end when the field approaches a minima with zero or negligible potential energy.

³⁹For a recent review, see [B. A. Bassett, S. Tsujikawa and D. Wands, Rev. Mod. Phys. 78, 537 \(2006\)](#).



Driving inflation with scalar fields




Inflation can be achieved with scalar fields encountered in high energy physics⁴⁰.

⁴⁰Image from P. J. Steinhardt, *Sci. Am.* **304**, 34 (2011).




A variety of potentials to choose from



Large_field

$$V(\phi) = \Lambda^4 (\phi/\mu)^p$$

$$V(\phi) = \Lambda^4 e^{\phi/\mu}$$



Small_field

$$V(\phi) = \Lambda^4 [1 - (\phi/\mu)^p]$$



Hybrid

$$V(\phi) = \Lambda^4 [1 + (\phi/\mu)^p]$$

A variety of scalar field potentials have been considered to drive inflation⁴¹.

⁴¹Image from W. Kinney, astro-ph/0301448.



The origin of the perturbations

- ♦ Inflation is typically driven with the aid of scalar fields. It is the quantum fluctuations associated with these scalar fields which are responsible for the origin of the perturbations.



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology
- 6 Useful references




The evolution of the perturbations

- ◆ The perturbations are amplified during the inflationary epoch, which leave their imprints as anisotropies in the CMB.
- ◆ The fluctuations in the CMB in turn grow in magnitude due to gravitational instability and develop into the structures that we see around us today.

[▶ Play movie](#)[▶ Play movie](#)

The timeline of the universe




A pictorial timeline of the universe⁴².

⁴²See http://wmap.gsfc.nasa.gov/media/060915/060915_CMB_Timeline150.jpg.



'Effects' of the cosmological parameters on the CMB⁴³





Sensitivity of the CMB angular power spectrum to the four cosmological parameters: Ω , Ω_A , $\Omega_B h^2$ and the non-relativistic matter density $\Omega_{NR} h^2$.

⁴³Figures from W. Hu and S. Dodelson, *Ann. Rev. Astron. Astrophys.* **40**, 171 (2002).



CMB anisotropies as seen by WMAP and Planck



Left: All-sky map of the anisotropies in the CMB created from nine years of Wilkinson Microwave Anisotropy Probe (WMAP) data⁴⁴.


Right: CMB intensity map derived from the joint analysis of Planck, WMAP, and 408 MHz observations⁴⁵. The above images show temperature variations (as color differences) of the order of $200^\circ \mu\text{K}$.

⁴⁴Image from <http://wmap.gsfc.nasa.gov/media/121238/index.html>.

⁴⁵P. A. R. Ade *et al.*, arXiv:1502.01582 [astro-ph.CO].



CMB angular power spectrum from Planck




The CMB TT angular power spectrum from the Planck 2018 data (red dots with error bars) and the best fit Λ CDM model with a power law primordial spectrum (solid blue curve)⁴⁶

⁴⁶Planck Collaboration (N. Aghanim *et al.*), *Astron. Astrophys.* **641**, A6 (2020).



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability




Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



Formation of structures due to gravitational instability

[▶ Play again](#)

Simulation illustrating the formation of structures due to gravitational instability⁴⁷.

⁴⁷Images from <http://cfcp.uchicago.edu/lss/group.html>.



The millennium simulation

- ◆ The Millennium Run used more than 10 billion particles to trace the evolution of the matter distribution in a cubic region of the universe over 2 billion light years on a side⁴⁷.
- ◆ It kept busy the principal supercomputer at the Max Planck Society's Supercomputing Centre in Garching, Germany for more than a month.

[▶ Play movie](#)

⁴⁷See <http://www.mpa-garching.mpg.de/galform/virgo/millennium/>.



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology**
- 6 Useful references





The standard model of cosmology

- ◆ The universe is homogeneous and isotropic at length scales of the order of 100 Mpc and above.
- ◆ Baryons, *i.e.* matter as we know it, contribute less than 5% to the total density of the universe today. Most of the matter today is, in fact, dark and dominated by dark energy. Dark energy and pressureless (*i.e.* cold) dark matter contribute about 70% and 25% to the the density today, respectively.
- ◆ The inflationary epoch magnifies the tiny fluctuations in the quantum fields present at the beginning of epoch into classical perturbations.
- ◆ These inhomogeneities leave their imprints as anisotropies in the CMB.
- ◆ Gravitational instability then takes over, and converts the tiny perturbations in the CMB into the large scale structures that we see around us today as galaxies and clusters of galaxies.



Ongoing and future missions



The **BICEP** (top left), **Euclid** (top right), **Square Kilometer Array** (bottom left) and the **Dark Energy Survey** (bottom right) missions are expected to provide unprecedented amount and quality of cosmological data that can help us unravel the mysteries of the universe.



Plan of the talk

- 1 A survey of the universe
- 2 The composition and evolution of the smooth universe
- 3 The inflationary paradigm and the origin of the perturbations
- 4 Evolution of the perturbations and their observable signatures
- 5 The standard model of cosmology
- 6 Useful references



Popular books

- ① S. Weinberg, *The First Three Minutes* (Bantam, New York, U.S.A., 1977).
- ② J. Silk, *The Big Bang* (W. H. Freeman, San Francisco, U.S.A., 1988).



Undergraduate level textbooks

- ① F. Shu, *The Physical Universe* (University Science Books, Mill Valley, California, U.S.A., 1982).
- ② A. Liddle, *An Introduction to Modern Cosmology* (John Wiley and Sons, Chichester, England, 2003).
- ③ P. Schneider, *Extragalactic Astronomy and Cosmology: An Introduction* (Springer, Berlin, Germany, 2006).
- ④ B. Ryden, *Introduction to Cosmology* (Cambridge University Press, Cambridge, England, 2017).



Thank you for your attention